

New Zealand Public Health Surveillance Report Volume 16 Issue 1

References

Increase in *Neisseria meningitidis* group W invasive disease

1. Australian Government Department of Health. 2017. Invasive meningococcal disease national surveillance report, with a focus on MenW. October 2017.
2. Public Health England. 2017. Invasive meningococcal disease in England: annual laboratory confirmed reports for epidemiological year 2016 to 2017.
3. CDC. 2017. Enhanced meningococcal disease surveillance report, 2016. [cited 29 January 2018]. Available from: <https://www.cdc.gov/meningococcal/downloads/NCIRD-EMS-Report.pdf>
4. Tsang RSW, Hoang L, Tyrrell GJ, et al. Increase in *Neisseria meningitidis* serogroup W invasive disease in Canada: 2009–2016. *Can Commun Dis Rep*. 2017; 43(7/8): 144-9.
5. Lopez L, Sherwood J. 2014. The epidemiology of meningococcal disease in New Zealand in 2013. Institute of Environmental Science and Research Ltd (ESR). Wellington, New Zealand.
6. CDC. Prevention and control of meningococcal disease: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*. 2013; 62(RR02): 1-22.
7. Australian Government Department of Health. 2018. Meningococcal W disease: information for health professionals. [cited 29 January 2018]. Available from: <http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-meningococcal-W-info-hp.htm>
8. Ladhani SN, Beebejaun K, Lucidarme J, et al. Increase in endemic *Neisseria meningitidis* capsular group W sequence type 11 complex associated with severe invasive disease in England and Wales. *Clin Infect Dis*. 2015; 60(4): 578-85.
9. Campbell H, Parikh S, Borrow R, et al. Presentation with gastrointestinal symptoms and high case fatality associated with group W meningococcal disease (MenW) in teenagers, England, July 2015 to January 2016. *Euro Surveill*. 2016; 21(12).
10. CDC. 2018. Meningococcal vaccination for adolescents: information for healthcare professionals. [cited 29 January 2018]. Available from: <https://www.cdc.gov/vaccines/vpd/mening/hcp/adolescent-vaccine.html>

Norovirus surveillance April 2016–December 2017: emergence of GII.P16-GII.4 Sydney_2012

1. Van Beek J, de Graaf M, Al-Hello H, et al. Molecular surveillance of norovirus, 2005–16: an epidemiological analysis of data collected from the NoroNet network. *Lancet Infect Dis*. 2018. [http://dx.doi.org/10.1016/S1473-3099\(18\)30059-8](http://dx.doi.org/10.1016/S1473-3099(18)30059-8)
2. Hewitt J. Norovirus outbreak surveillance - a new GII.4 variant emerges in 2012. *New Zealand Public Health Surveillance Report*. 2013; 11(1): 8. https://surv.esr.cri.nz/PDF_surveillance/NZPHSR/2013/NZPHSRMar2013.pdf
3. Barreira D, Fumian TM, Tonini MAL, et al. Detection and molecular characterization of the novel recombinant norovirus GII.P16-GII.4 Sydney in southeastern Brazil in 2016. *PLoS One*. 2017; 12(12): e0189504. <https://doi.org/10.1371/journal.pone.0189504>

4. Bidalot M, Théry L, Kaplon J, et al. Emergence of new recombinant noroviruses GII.p16-GII.4 and GII.p16-GII.2, France, winter 2016 to 2017. *Euro Surveill.* 2017; 22(15): 30508. <http://dx.doi.org/10.2807/1560-7917.ES.2017.22.15.30508>
5. Cannon JL, Barclay L, Collins NR, et al. Genetic and epidemiologic trends of norovirus outbreaks in the United States from 2013 to 2016 demonstrated emergence of novel GII.4 recombinant viruses. Tang Y-W, ed. *J Clin Microbiol.* 2017; 55(7): 2208-21.
6. Choi YS, Koo ES, Kim MS, et al. Re-emergence of a GII.4 norovirus Sydney 2012 variant equipped with GII.P16 RdRp and its predominance over novel variants of GII.17 in South Korea in 2016. *Food Environ Virol.* 2017; 9(2): 168-78.
7. White P, Lun J, Hewitt J, et al. Emerging recombinant noroviruses identified by clinical and wastewater screening. *Emerg Microbes Infect.* 2018. In press.
8. Lindesmith LC, Brewer-Jensen PD, Mallory ML, et al. Antigenic characterization of a novel recombinant GII.P16-GII.4 Sydney norovirus strain with minor sequence variation leading to antibody escape. *J Infect Dis.* 2017. <https://doi.org/10.1093/infdis/jix651>
9. Ruis C, Roy S, Brown JR, et al. The emerging GII.P16-GII.4 Sydney 2012 norovirus lineage is circulating worldwide, arose by late-2014 and contains polymerase changes that may increase virus transmission. *PLoS One.* 2017; 12(6): e0179572. <https://doi.org/10.1371/journal.pone.0179572>