

National Wastewater Surveillance Programme | SARS-CoV-2

Week 30 (Week Ending 31 July 2022)

Report prepared on 03 August 2022

**SARS-CoV-2 RNA
 detected at
 97% (102/105)
 sites tested**


75%
 of NZ population
 covered by
 wastewater testing

**Most prevalent
 variant detected
 Omicron BA.4/5
 (~90%)**

**Overall, SARS-CoV-2 levels in wastewater have plateaued but
 there are site specific increases and decreases
 Omicron BA.4/5 levels continue to increase relative to BA.2**

SARS-CoV-2 in Wastewater and Reported COVID-19 Case Numbers

National, last 6 months, on log₁₀ scale



- Week ending **31 July 2022**, **35%** of sites have **increased** SARS-CoV-2 levels compared to the previous week, and **39% decreased** levels. There is a similar mixed trend compared to the previous month.

Results For Week 30 2022

In the week ending 31 July 2022, 156 samples were collected from 105 locations in New Zealand.

SARS CoV-2 RNA was detected from 102 (97%) of the sites tested this week (Figure 1, Figure 2 and Table 1). SARS-CoV-2 RNA was **not detected** at three sites: Reefton (collected 25/7/2022), Kaponga (collected 26/7/2022), and Porangahau (collected 29/7/2022). A further two sites tested positive for SARS-CoV-2 RNA in only one of the two samples collected for the week: Woodville (negative on 26/7, positive on 27/7) and Kaiwaka (positive on 25/7, negative on 28/7).

Figure 1 below gives the proportion of sites with positive results for this and previous weeks.




Figure 1. Results for SARS-CoV-2 RNA in wastewater collected across New Zealand.

Using a threshold of an increase of 30% or more (Figure 2), 35% of sites have higher levels compared with 4 weeks ago.




Figure 2. Comparison of SARS-CoV-2 levels for the week ending 31 July 2022, compared with the levels measured **1 week, 2 weeks, 4 weeks, or 12 weeks previously**. Only sites with results for both time points are included. When the viral quantity is 30% or more higher this is labelled as increased (red up arrow on map). When the viral quantity is 30% or more lower this is labelled as decreased (blue down arrow on map). If viral levels have changed less than this in the compared weeks, this is labelled as no change (yellow circle on map).

Regional summaries (Figure 3) also indicate a **mixed trend, with decreases in levels** of genome copies per person per day (rolling 2-week simple average) across Northern, Central and Southern regions and **increases** in Auckland Metro and Te Manawa Taki regions compared with the previous week.




Figure 3. Regional trends in SARS-CoV-2 levels in wastewater. Northern Region excludes the Auckland metro sites (Eastern, Western, Southwestern and Rosedale).

Consistent with whole genome sequencing (WGS) of clinical cases, **Omicron BA.4/5 variants are now firmly established as the most commonly circulating variants across Aotearoa (national average of ~90%)**. Over the past 7 weeks nationally (aggregated across all 21 sentinel sites), BA.4/5 has increased from less than 5%, to approximately 90% (Figure 4B).

Following the BA.1 detection last week in Wellington (Moa Point), this week Hutt Valley is the only site where this variant is detected (Figure 4A).





Figure 4. Variants present in New Zealand wastewater between 12 June and 31 July 2022, based on data from up to 22 sentinel sites surveyed each week. The level of precision and sensitivity in these percentage estimates can be uncertain.

Interpreting site graphs

Interpreting Sites Graphs




- 1** Site Name
- 2** Sample Collection method and population. Results based on autosampler may be more representative than grab sample-based results.
- 3** Wastewater Results Shown as Solid Line | Rolling 14-day average of genome copies/person/day on a log₁₀ scale.
- 4** Individual Results Samples Shown as Circles | Rolling 14-day average of genome copies/person/day on a log₁₀ scale.
- 5** Rolling 7-day average new cases shown as dashed line | New cases reported in a catchment based on reported date of illness on a log₁₀ scale. This data is not available for all sites & subject to change.


Note: Wastewater and cases data are on a log₁₀ scale. Scales on all graphs have been normalised to cover the same scale on every graph. Care should be taken when interpreting the data.

Northland


Whangarei Autosampler 65.0K




Dargaville Grab 5.0K





Hikurangi Grab 1.7



Ruakaka Grab 4.5K




Mangawhai Grab 1.1K




Status ● Detected ● Not detected

Auckland


Auckland (Combined) Autosampler 1.1M




North Shore Autosampler 240.0K




Auckland East Autosampler 680.0K




Auckland Southwest Autosampler 120.0K



Auckland West Autosampler 315.0K




Army Bay Autosampler 42.0K




Status ● Detected ● Not detected


Pukekohe *Grab 20.9K*




Snells/Algies *Autosampler 4.0K*




Clarks Beach *Grab 2.0K*




Waiuku *Grab 7.9K*




Helensville *Autosampler 3.8K*




Wellsford *Autosampler 1.7K*




Beachlands *Grab 6.8K*



Warkworth *Autosampler 3.5K*




Omaha *Autosampler 1000*




Cases - 7 day rolling average

Waikato


Hamilton Autosampler 169.0K





Taupo Autosampler/Grab 23.0K



Cambridge Autosampler 20.1K




Te Awamutu Autosampler 13.1K




Status ● Detected ● Not detected

Cases - 7 day rolling average


Thames Autosampler 7.5K




Whitianga Autosampler 6.6K




Whangamata Autosampler 4.0K





Bay of Plenty and Gisborne
Mt Maunganui Autosampler 65.0K




Gisborne Autosampler 37.0K




Rotorua Autosampler 59.0K



Whakatane Autosampler 21.0K




Tauranga Autosampler 50.0K




Status ● Detected ● Not detected


Te Puke Autosampler 9.7K




Kawerau Autosampler 7.0K




Waihi Beach Autosampler 3.6K




Katikati Autosampler 5.5K



Maketu Autosampler 1.3K




Opotiki Grab 3.8K




Cases - 7 day rolling average

Hawke's Bay


Hastings Autosampler 80.0K




Wairoa Grab 4.4K




Napier Autosampler 55.0K



Waipawa Autosampler 2.2K



Waipukurau Autosampler 4.6K




Status ● Detected ● Not detected


Cases - 7 day rolling average

Taranaki


New Plymouth Autosampler 88.0K




Eltham Autosampler 2.0K



Hawera Autosampler 12.0K




Opunake Autosampler 1.4K




Status ● Detected ● Not detected

Patea Autosampler 1.2K




Manawatu-Whanganui


Palmerston North Autosampler 90.0K




Whanganui Urban Autosampler 44.5K




Levin Autosampler 21.2K



Dannevirke Grab 5.7K




Taumarunui Grab 4.0K




Status ● Detected ● Not detected


Pahiatua *Grab 2.8K*



Woodville *Grab 1.7K*




Eketahuna *Grab 1.6K*




Cases - 7 day rolling average

Wellington


Wellington (Moa Point) Autosampler 168.0K




Paraparaumu Autosampler 49.0K




Hutt Valley Autosampler 133.0K




Masterton Grab 20.7K



Porirua Autosampler 85.0K




Wellington (Western) Autosampler 14.0K




Status ● Detected ● Not detected

Cases - 7 day rolling average


Carterton *Grab 5.8K*




Greytown *Grab 2.4K*




Otaki *Grab 3.5K*



Martinborough *Autosampler/grab 1.6K*




Featherston *Grab 2.5K*




SARS-CoV-2 genome copies / day / person

Cases - 7 day rolling average


Tasman, Nelson, and Marlborough
Richmond/Nelson South Autosampler 60.0K




Motueka Autosampler 8.3K




Blenheim Autosampler 31.0K



Picton Autosampler 5.0K




Nelson Central/North Autosampler 26.0K




Status ● Detected ● Not detected

West Coast and Canterbury


Christchurch Autosampler 368.0K




Rangiora Grab 19.0K




Rolleston & Eastern Selwyn Autosampler 35.0K





Ashburton Autosampler 18.0K



Timaru Autosampler 28.0K




Kaipoi Grab 12.5K




Status ● Detected ● Not detected


Oamaru Autosampler 12.0K




Westport Grab 5.0K




Kaikoura Grab 2.4K




Greymouth Grab 10.0K



Leeston Autosampler 3.9K




Amberley Grab 1.8K




Cases - 7 day rolling average


Woodend Grab 7.6K



Hokitika Grab 2.9K




Reefton Grab 1000




Otago and Southland



Dunedin (Tahuna) Autosampler 84.0K




Dunedin (Green Island) Autosampler 22.9K




Invercargill Autosampler 50.0K



Dunedin (Mosgiel) Autosampler 14.6K




Queenstown Autosampler 40.0K




Status ● Detected ● Not detected


Wanaka Grab 14.5K




Gore Autosampler 8.0K




Balclutha Grab 4.1K




Cromwell Autosampler 7.1K



Bluff Autosampler 2.0K



Alexandra Autosampler 6.2K



Cases - 7 day rolling average

Table 1. Weekly summary

Key: ■ Not Detected ■ Detected (below limit of quantification) ■ Detected (quantifiable) ■ Not sampled.

Site	2021																2022																														
	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Northland																																															
Ahipara		■			■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■																					
Awanui		■																																													
Dargaville		■							■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Haruru																				■	■	■	■	■	■	■	■																				
Hihi		■																																													
Hikurangi		■																																													
Kaeo		■																																													
Kaikohe		■																																													
Kaitaia	■																			■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■		
Kaiwaka	■																																														
Kawakawa	■																																														
Kerikeri		■																																													
Kohukohu		■																																													
Mangawhai		■																																													
Maungaturoto		■																																													
Opononi		■																																													
Paihia		■																																													
Rangiputa		■																																													
Rawene		■																																													
Ruakaka	■																																														
Russell		■																																													
Taipa		■																																													
Waipu	■																																														
Whangarei	■																																														
Whatuwhiwhi	■																																														
Auckland																																															
Army Bay		■																																													
Auckland (Combined)																																															
Auckland East																																															
Auckland Southwest																																															
Auckland West																																															
Beachlands		■																																													
Clarks Beach		■																																													
Helensville		■																																													
Kawakawa Bay		■																																													
Kingseat		■																																													
North Shore		■																																													
Omaha		■																																													
Pukekohe	■																																														
Snells/Algies		■																																													
Waiheke Island		■																																													
Waiuku		■																																													

Acknowledgements

This work represents the combined efforts of a large number of individuals and organisations.

We continue to be indebted to the teams across the country who are collecting the wastewater that underpins this work.

The wastewater analysis has been undertaken at the ESR Kenepuru (in Porirua) and Christchurch laboratories by a team which may on any given week include contributions from: Kirsten Carter, Joanne Chapman, Dawn Croucher, Joanne Hewitt, Joycelyn Ho, Anower Javed, Susan Lin, Olivia Macrae, Ashley McDonald, Andrew Ng, Ashley Orton, Paula Scholes and Fatiha Sulthana. Data science analysis, visualisation and reporting is the result of team effort from: Franco Andrews, Bridget Armstrong, Raewyn Campbell, Gerhard de Beer, Richard Dean, Brent Gilpin, Joanne Hewitt, Dawen Li, Helen Morris and Bindu Priya. Ongoing support for this work from the Ministry of Health and ESR management is appreciated.

Notes

Overview

SARS-CoV-2, the virus that causes COVID-19 disease, is shed in the faeces of people that are infected and so the viral RNA can be detected in wastewater. As such, testing wastewater for SARS-CoV-2 RNA is an efficient population-based COVID-19 surveillance tool. Based on national and international data, this method has been shown to be an indicator of increasing and decreasing cases (i.e., early warning system) and complements other surveillance tools. A national wastewater COVID-19 surveillance programme was established in 2021 by the Institute of Environmental Science and Research (ESR). This work is funded by the New Zealand Ministry of Health and is part of New Zealand's COVID-19 response.

Wastewater samples are collected from wastewater treatment plants across both the North and South Island of New Zealand. Grab or 24 hr composite samples are collected. Most sites are sampled at least weekly between Monday and Thursday of any given week. The number of sites and frequency of collection varies over time. Greater variability is expected with grab samples.

Approach

Samples are sent from each wastewater treatment plant to one of the ESR laboratories (Porirua or Christchurch). Processing involves the concentration of virus and extraction of viral RNA. The presence of SARS-CoV-2 RNA in the sample is then determined using RT-qPCR.

A result of **not detected**, means that SARS-CoV-2 RNA is either absent from the sample, or at a level too low to be detected. When SARS-CoV-2 RNA is **detected**, the concentration in the sample can be calculated. Low amounts of SARS-CoV-2 RNA in a sample may not be able to be accurately quantified and are recorded as less than the limit of quantitation. For quantitation, the raw concentration data (i.e., genome copies per L) is converted to a viral load of **genome copies per day per person**. This calculation considers the flow rate of wastewater entering the wastewater treatment plant and the population in the catchment. This is the population-normalised viral load.

Sites and frequency of sample collection: The catchment population sites selected for the surveillance range from approximately 100 to over 1,000,000 individuals. The sites cover all regions of the country. Most major towns and all cities, as well as many smaller communities, are included. In early 2022, the wastewater catchment areas cover over 80% of the population connected to wastewater treatment plants. The sites from which samples have been collected have varied over the last 12 months. New sites may be added over

time, and/or sampling may reduce in frequency or cease for other sites. The selection and frequency of sampling vary depending on the local population, access to wastewater collection points, staff availability to collect samples and risk factors. When included, samples are collected at least weekly, with twice weekly sampling being common. A number of samples have also been collected from non-WWTP sites (manholes and pump stations- mostly in Auckland).

Sampling method: The preferred option is to automatically collect a 24 hour 'composite' sample. This is where a pump automatically collects a small volume of wastewater every 15 minutes over 24 hours using a composite sampler. These samplers are available in some wastewater treatment plants. When composite samplers are not available, 'grab' samples are collected. These range from a sample being taken at a single point in time, to 3 samples taken over 30 minutes, to samples collected over a day. Grab samples represent only the composition of the source at that time of collection and may not be as representative as a 24-hour composite sampler. More variation may be expected with grab samples.

Laboratory analysis of wastewater samples: Samples are sent from each wastewater treatment plant to one of the ESR laboratories (Porirua or Christchurch). Processing of each sample commences within an hour or two of receipt. Processing involves the concentration of virus from 250 mL sample to approx. 1 mL using centrifugation and polyethylene glycol. Viral RNA is then extracted from a small volume of 0.2 mL concentrate to give a final volume of 0.05 mL. The presence of SARS-CoV-2 RNA is determined using RT-qPCR. SARS-CoV-2 is considered detected when any of the RT-qPCR replicates are positive.

RT-qPCR: Reverse transcription (RT) is used to convert RNA to complementary DNA (cDNA), followed by quantitative PCR (qPCR) to amplify and quantify the cDNA. RT-qPCR is the standard method used for detection and quantification of viral RNA

Method sensitivity: The protocol used to concentrate SARS-CoV-2 from wastewater allows for the sensitive detection of SARS-CoV-2 by RT-qPCR. ESR has shown that when 10 individuals are actively shedding SARS-CoV-2 RNA in a catchment of 100,000 individuals, there was a high likelihood of detecting viral RNA in wastewater (<https://doi.org/10.1016/j.watres.2021.118032>). Shedding by one individual may be detected in wastewater, but it does depend on many factors including the amount and duration of shedding. Very low levels in wastewater may be not able to be quantified (i.e., less than the limit of quantification- see below).

SARS-CoV-2 RNA detected (positive result): A positive detection in the wastewater indicates that at least one person has been shedding SARS-CoV-2 into the wastewater at some point during the time period that the sample was being collected. In some cases, detections could also be due to the shedding of low levels of SARS-CoV-2 RNA by a recently recovered case. The detection of SARS-CoV-2 RNA does not indicate that infectious virus is present.

SARS-CoV-2 RNA not detected (negative result): A negative result can occur because there are no active 'shedding' cases in the catchment or because the SARS-CoV-2 RNA concentration is too low to be detected, most likely because there are a very low number of cases in the wastewater catchment. Therefore, negative finding does not necessarily guarantee the absence of COVID-19 in the community.

Viral loads and normalisation: When detected, the SARS-CoV-2 RNA concentration is calculated as genome copies per L of wastewater. This is then converted to a viral load of **genome copies/day/person**. This conversion takes into account the flow rate of wastewater entering the treatment plant (the influent) and the population in the catchment. The **flow rate** is the total volume (m³ per day) recorded at the inlet of the wastewater treatment plant over 24 hours. This is a **population-normalised viral load**. Currently, the flow

rate is the average annual flow rate, but will be replaced with daily flow rate when available (note that rainfall may significantly increase the flow rate at the inlet, diluting the sample, and may result in lower concentrations and a false negative result).

In future, SARS-CoV-2 RNA concentrations will also be normalised by testing for the presence of pepper mild mottled virus (PMMoV). PMMoV is a virus that infects peppers but not humans. Consumption of peppers or pepper products, such as chilli sauce, means that PMMoV is detected in wastewater – normally at very high concentrations. Therefore, PMMoV has been found to be a useful proxy for the amount of faecal material in a wastewater sample. For normalisation, the concentration of SARS-CoV-2 RNA is divided by that of PMMoV in each sample. Different normalisation methods may result in changes to some data points, but trends are unlikely to change significantly.

Limit of quantification: The lowest concentration of the target that can be reliably quantified is referred to as the limit of quantification. For those samples where SARS-CoV-2 is detected but cannot be quantified, a value of 5 genome copies/mL wastewater is used. While a standard method is being used, virus recovery can vary from sample to sample, and this may affect the quantitation.

Data subject to change: Data generated for the New Zealand Wastewater COVID-19 Surveillance Programme should be considered provisional and may be subject to change. Data may be incomplete for the most recent 2-week period due to processing, testing and reporting delays.

Data not shown:

- Data from 'ad hoc' sampling locations including from individual facilities/building (e.g., workplaces, prisons, MIQs) are not included.
- Results from certain samples may not be shown, as the result was either deemed invalid, or the sample could not be tested (e.g., leaked in transit, not labelled).

Key Points & Limitations

- SARS-CoV-2 RNA concentrations should not be compared between wastewater catchments.
- Day to day variability in SARS-CoV-2 RNA concentrations, especially in smaller catchments, is to be expected. Greater variability is expected with grab samples.
- Generally, increasing viral loads are associated with increasing numbers of people with SARS-CoV-2 infection and vice versa (decreasing concentrations indicating decreasing cases). However, there are a number of factors that affect the amount of viral RNA detected and so data from wastewater surveillance cannot indicate the exact number of COVID-19 cases in the catchment area.
- The number of COVID-19 cases reported via individual testing are reported for each region to provide a comparison to the wastewater results. The cases in each catchment area are an estimate of the number of people in that wastewater catchment area that have reported a positive test. However, because the wastewater catchments do not exactly align with regional boundaries, the number of cases estimated by region and by water catchment area may be different.
- Data are provisional and may be subject to change by location.
- As septic tank systems are not connected to wastewater treatment plants, the wastewater from these households will not be represented in the data.

For further information please contact

Brent Gilpin
Science Leader
Brent.gilpin@esr.cri.nz

Joanne Hewitt
Senior Scientist, Environmental & Food Virology
Joanne.hewitt@esr.cri.nz

END OF REPORT